


1 (currently amended). A compound of the formula I:



wherein

A is a 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, and 12-membered mono-, bi- or spirobicyclic ring containing one or more heteroatoms selected from the group of N, O and S, and is optionally substituted with F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)-alkyl, aryl, CON(R11)(R12), N(R13)(R14), OH, O-(C₁-C₆)-alkyl, S-(C₁-C₆)-alkyl, N(R15)CO(C₁-C₆)-alkyl or COO-(C₁-C₆)-alkyl;

R11, R12, R13, R14, R15 are each independently H, (C₁-C₆)-alkyl or a heterocycle;

n is 1;

m is 0, 1, 2, 3, 4, 5 or 6;

R1 is R8, (C₁-C₆)-alkylene-R8, (C₂-C₆)-alkenylene-R9, (SO₂)-R8, (SO₂)-(C₁-C₆)-alkylene-R8, (SO₂)-(C₂-C₆)-alkenylene-R9, (C=O)-R8, (C=O)-(C₁-C₆)-alkylene-R8, (C=O)NH-R8, (C=O)-(C₂-C₆)-alkenylene-R9, (C=O)-NH-(C₁-C₆)-alkylene-R8, (C=O)-NH-(C₂-C₆)-alkenylene-R9, COO-R8, COO-(C₁-C₆)-alkylene-R8, COO-(C₂-

C₆)-alkenylene-R₉, alkynylene-R₉ or (C₁-C₄-alkyl)-heterocycle, wherein the alkylene component of said (C₁-C₆)-alkylene-R₈, (C₂-C₆)-alkenylene-R₉, (SO₂)-(C₁-C₆)-alkylene-R₈, (SO₂)-(C₂-C₆)-alkenylene-R₉, (C=O)-(C₁-C₆)-alkylene-R₈, (C=O)-(C₂-C₆)-alkenylene-R₉, (C=O)-NH-(C₁-C₆)-alkylene-R₈, (C=O)-NH-(C₂-C₆)-alkenylene-R₉, COO-(C₁-C₆)-alkylene-R₈, COO-(C₂-C₆)-alkenylene-R₉ and alkynylene-R₉ groups is optionally substituted by F;

- R₈, R₉ are each independently H, F, Cl, Br, I, OH, CF₃, aryl, heterocycle or (C₃-C₈)-cycloalkyl, wherein said aryl, heterocycle and (C₃-C₈)-cycloalkyl groups are optionally mono-, di- or tri-substituted by F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, NH₂, CON(R₁₁)(R₁₂), N(R₁₃)(R₁₄), SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CONH₂;
- R₂ is NH₂, NO₂, N(R₁₃)(R₁₄), NH-SO₂-CH₃, NH-SO₂-R₁₂, NR₁₁-SO₂-R₁₂, N(CO)R₁₁, NHCONR₁₁, N(C₁-C₆-alkyl)N⁺(C₁-C₄-alkyl)₃ or a nitrogen-containing heterocycle, wherein said heterocycle is bonded via a nitrogen atom;
- R₃, R₄, R₅ are each independently H, F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, O-(C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, S-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₃-C₈)-cycloalkyl, O-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkenyl, O-(C₃-C₈)-cycloalkenyl, (C₂-C₆)-alkynyl, aryl, O-aryl (C₁-C₈)-alkylene-aryl, O-(C₁-C₈)-alkylene-aryl, S-aryl, N((C₁-C₆)-alkyl)₂, SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CO-N((C₁-C₆)-alkyl)₂;
- R₆ is H, F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, O-(C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, S-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₃-C₈)-cycloalkyl, O-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkenyl, O-(C₃-C₈)-cycloalkenyl, (C₂-C₆)-alkynyl, (C₀-C₈)-alkylene-aryl, O-(C₀-C₈)-alkylene-aryl, S-aryl, N((C₁-C₆)-alkyl)₂, SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CO-N((C₁-C₆)-alkyl)₂;

Aryl is phenyl or naphthyl;

Heterocycle is acridinyl, azocinyl, benzimidazolyl, benzofuryl, benzothienyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxalinyl, quinuclidinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]-tetrahydrofuran, furyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, 1H-indazolyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, benzimidazolyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazoles, pyridoimidazoles, pyridothiazoles, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, thiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thienyl, triazolyl, tetrazolyl and xanthenyl and corresponding N-oxides;


wherein said heterocycle is optionally substituted one or more times, each substituent independently chosen from F, Cl, Br, I, CF₃, NO₂, N₃, CN, COOH, COO(C₁-C₆)alkyl, CONH₂, CONH(C₁-C₆)alkyl, CON[(C₁-C₆)alkyl]₂, (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, O-(C₁-C₆)-alkyl, wherein one or more than one, or all hydrogen(s) in the alkyl radicals may be replaced by fluorine;

PO₃H₂, SO₃H, SO₂-NH₂, SO₂NH(C₁-C₆)-alkyl, SO₂N[(C₁-C₆)-alkyl]₂, S-(C₁-C₆)-alkyl, S-(CH₂)_n-phenyl, SO-(C₁-C₆)-

alkyl, SO-(CH₂)_n-phenyl, SO₂-(C₁-C₆)-alkyl, SO₂-(CH₂)_n-phenyl, wherein n can be 0-6, and the phenyl radical may be substituted up to two times by F, Cl, Br, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl or NH₂; C(NH)(NH₂), NH₂, NH-(C₁-C₆)-alkyl, N((C₁-C₆)-alkyl)₂, NH(C₁-C₇)-acyl, phenyl and O-(CH₂)_n-phenyl, wherein n may be 0-6, and wherein the phenyl ring may be substituted one to 3 times by F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, NH₂, NH(C₁-C₆)-alkyl, N((C₁-C₆)-alkyl)₂, SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CONH₂;

and pharmaceutically acceptable salts thereof.

2 (original). The compound of Claim 1 having the following structure Ia



wherein

A is a 3-, 4-, 5-, 6-, 7-, 8-, 9-, 10-, 11-, and 12-membered mono-, bi- or spirobicyclic ring containing one or more heteroatoms selected from the group of N, O and S, and is optionally substituted with F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)-alkyl, aryl,

CON(R11)(R12), N(R13)(R14), OH, O-(C₁-C₆)-alkyl, S-(C₁-C₆)-alkyl, N(R15)CO(C₁-C₆)-alkyl or COO-(C₁-C₆)-alkyl;

R11, R12, R13, R14, R15 are each independently H, (C₁-C₆)-alkyl or a heterocycle;

m is 0, 1, 2, 3, 4, 5 or 6;

R1 is R8, (C₁-C₆)-alkylene-R8, (C₂-C₆)-alkenylene-R9, (SO₂)-R8, (SO₂)-(C₁-C₆)-alkylene-R8, (SO₂)-(C₂-C₆)-alkenylene-R9, (C=O)-R8, (C=O)-(C₁-C₆)-alkylene-R8, (C=O)NH-R8, (C=O)-(C₂-C₆)-alkenylene-R9, (C=O)-NH-(C₁-C₆)-alkylene-R8, (C=O)-NH-(C₂-C₆)-alkenylene-R9, COO-R8, COO-(C₁-C₆)-alkylene-R8, COO-(C₂-C₆)-alkenylene-R9, alkynylene-R9 or (C₁-C₄-alkyl)-heterocycle;

R8, R9 are each independently H, F, Cl, Br, I, OH, CF₃, aryl, heterocycle or (C₃-C₈)-cycloalkyl, wherein said aryl, heterocycle and (C₃-C₈)-cycloalkyl groups are optionally mono-, di- or tri-substituted by F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, NH₂, CON(R11)(R12), N(R13)(R14), SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CONH₂;

R2 is NH₂, NO₂, N(R13)(R14), NH-SO₂-CH₃, NH-SO₂-R12, NR11-SO₂-R12, N(CO)R11, NHCONR11, N(C₁-C₆-alkyl)N⁺(C₁-C₄-alkyl)₃ or a nitrogen-containing heterocycle, wherein said heterocycle is bonded via a nitrogen atom;

R3, R4, R5 are each independently H, F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, O-(C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, S-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₃-C₈)-cycloalkyl, O-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkenyl, O-(C₃-C₈)-cycloalkenyl, (C₂-C₆)-alkynyl, aryl, O-aryl (C₁-C₈)-alkylene-aryl, O-(C₁-C₈)-alkylene-aryl, S-aryl, N((C₁-C₆)-alkyl)₂, SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CO-N((C₁-C₆)-alkyl)₂;

R6 is H, F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, O-(C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, S-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₃-C₈)-cycloalkyl, O-(C₃-

C₈)-cycloalkyl, (C₃-C₈)-cycloalkenyl, O-(C₃-C₈)-cycloalkenyl, (C₂-C₆)-alkynyl, aryl, O-aryl, (C₁-C₈)-alkylene-aryl, O-(C₁-C₈)-alkylene-aryl, S-aryl, N((C₁-C₆)-alkyl)₂, SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CO-N((C₁-C₆)-alkyl)₂;

and pharmaceutically acceptable salts thereof.

3 (original). The compound of Claim 2 wherein

A is aryl wherein said aryl is optionally substituted by F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)-alkyl, aryl, CON(R11)(R12), N(R13)(R14), OH, O-(C₁-C₆)-alkyl, S-(C₁-C₆)-alkyl, N(R15)CO(C₁-C₆)-alkyl or COO-(C₁-C₆)-alkyl;

R11, R12, R13, R14, R15 are each independently H, (C₁-C₆)-alkyl or heterocycle;

m is 1;

R1 is R8, (C₁-C₆)-alkylene-R8, (C₂-C₆)-alkenylene-R9, (SO₂)-R8, (SO₂)-(C₁-C₆)-alkylene-R8, (SO₂)-(C₂-C₆)-alkenylene-R9, (C=O)-R8, (C=O)-(C₁-C₆)-alkylene-R8, (C=O)NH-R8, (C=O)-(C₂-C₆)-alkenylene-R9, (C=O)-NH-(C₁-C₆)-alkylene-R8, (C=O)-NH-(C₂-C₆)-alkenylene-R9, COO-R8, COO-(C₁-C₆)-alkylene-R8, COO-(C₂-C₆)-alkenylene-R9, alkynylene-R9 or (C₁-C₄-alkyl)-heterocycle;

R8, R9 are each independently H, F, Cl, Br, I, OH, CF₃, aryl, heterocycle or (C₃-C₈)-cycloalkyl, wherein said aryl, heterocycle and (C₃-C₈)-cycloalkyl groups are optionally mono-, di-, or tri-substituted by F, Cl, Br, I, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)-alkyl, (C₁-C₆)-alkyl, NH₂, CON(R11)(R12), N(R13)(R14), SO₂-CH₃, COOH, COO-(C₁-C₆)-alkyl or CONH₂;

R2 is NH₂, NO₂, N(R13)(R14), NH-SO₂-CH₃, NH-SO₂-R12, NR11-SO₂-R12, N(CO)R11, NHCONR11, N(C₁-C₆-alkyl)N⁺(C₁-C₄-alkyl)₃ or a nitrogen-containing heterocycle, wherein said heterocycle is bonded via a nitrogen atom,

R3 is H

R4, R5 are each independently H, F, Cl, Br, OH, CF₃, OCF₃, O-(C₁-C₆)-alkyl or (C₁-C₆)-alkyl;

R6 is H;

and pharmaceutically acceptable salts thereof.

4 (original). The compound of Claim 3 wherein

A is aryl, wherein said aryl group is optionally substituted by F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)-alkyl, aryl, CON(R11)(R12), N(R13)(R14), OH, O-(C₁-C₆)-alkyl, S-(C₁-C₆)-alkyl, N(R15)CO(C₁-C₆)-alkyl or COO-(C₁-C₆)-alkyl;

R11, R12, R13, R14, R15 are each independently H, (C₁-C₆)-alkyl or heterocycle;

m is 1;

R1 is (C₁-C₆)-alkyl or (C₁-C₆)-alkylene-R8;

R8, R9 are each independently F, Cl, Br, I, OH or CF₃;

R2 is NH₂, NO₂, CN, N(R13)(R14), NH-SO₂-CH₃, NH-SO₂-R12, NR11-SO₂-R12, N(CO)R11, NHCONR11, N(C₁-C₆-alkyl)N⁺(C₁-C₄-alkyl)₃ or a nitrogen-containing heterocycle, wherein said heterocycle is bonded via a nitrogen atom,

R3 is H;

R4 is F, Cl, Br, OH, CF₃, OCF₃, O-(C₁-C₆)-alkyl or (C₁-C₆)-alkyl;

R5 is H, F, Cl, Br, OH, CF₃, OCF₃, O-(C₁-C₆)-alkyl or (C₁-C₆)-alkyl;

R6 is H;

and pharmaceutically acceptable salts thereof.

5 (original). A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

6-8 (canceled).

9 (original). A method of treating obesity comprising administering to a patient in need thereof a compound of Claim 1.

10-12 (canceled).

13 (original). A method of reducing weight in mammals comprising administering to a patient in need thereof a compound of Claim 1.

14-15 (canceled).